



Quality and Safety of Locally Produced Chicken Luncheon

Doaa M.A. Abdel-Allah, Heba H.S. Abdel-Naeem and Amal M.A. El-Sherif
Department of Food Hygiene and Control, Faculty of Veterinary Medicine, Cairo University

Abstract

Ninety samples of Egyptian chicken luncheon sausage produced by six different meat processing plants (Fifteen samples each) were collected from different production lots within one month after processing from the supermarkets in Giza and Cairo. These samples were subjected to sensory investigation, determination of deterioration criteria, bacteriological examinations as well as detection of non-meat tissues by histological examination to assure their quality and safety. The obtained results revealed that all samples had deteriorated sensory attributes, high TBA, pH value as well as very high bacteriological load. The results of histological examination showed that low skeletal muscular tissues and high fat content were evident in all samples. Moreover, bone particles, cartilage particles, skin tissue with feather and other non meat tissue such as digestive tract were observed.

Keywords: Chicken/Luncheon sausage/ pH/TBA/Bone/Cartilage/Skin.

Introduction

Poultry meat production and consumption has dramatically increased over the last few decades (American Meat Institute, 2004). This increase can be attributed to low production costs, rapid growth rates, high nutritional values, and the increased production of further processed products (Barbut, 2002). For several reasons, people prefer poultry meat products as compared to beef or pork. The manufacture of such products usually costs less than that of similar beef and pork products (Guerrero-Legarreta and Hui, 2010). An added benefit is that poultry meat is not restricted by most cultural and religious laws, and it is consumed by both Jews and Muslims (Deumier and Collignan,

The first consumer right is to have a product of good quality and not constituting any health hazard. Quality products are those that meet some need or expectation of consumers and are safe and wholesome as well. The acceptance of further processed chicken meat products depends upon overall-acceptability, The increasing demand for meat products as well as the relative shortage in raw meat has led to its adulteration with various tissues and non-meat ingredients (Emara and Nouman, 2002). Mechanically deboned meat is commonly used as a major ingredient in producing of emulsified and comminuted meat products such as chicken sausages, frankfurters and burger either as a binding agent or as an inexpensive source of

2003). Further processing of poultry meat involves conversion of raw poultry carcasses into value added products (AL-Dughaym and Altabari, 2010). The production and consumption of chicken luncheon sausages has been increasing globally in particular chicken emulsion sausage (Bonoli et al., 2007 and Hwang et al., 2011). These luncheon sausages are becoming more popular due to their sensory characteristics and ease of preparation, which reflects the development of more functionality enhanced chicken emulsion sausages with added dietary fiber (Park et al., 2012). Emulsion based product is one such technology which can convert meat of low organoleptic value into a highly acceptable product without tenderization (Ilayabharathi et al., 2012).

color, odor, taste and consistency. Therefore, consumers had given much greater choice over the foods which are more selective, of high quality and cheap. Finally, the product quality became more significant factor in meat products marketing (Potter, 2001 and Agamy and Hegazy, 2011).

meat to reduce the cost of the products. Moreover, it offers the food industry with a raw material with excellent nutritional (high protein content), good technological and functional properties (Daros et al., 2005 and Püssa et al., 2009). However, it has some disadvantages, such as rapid onset of oxidative rancidity resulting in off-flavours and off-odours, colour and texture problems (Freitas et al., 2004) and

the high microbial load, which makes it a highly perishable raw material (Gill, 1988). Based on E.S.S.(1696/2005) view point, the use of undesirable organs of slaughter animals, including the visceral organs, hyaline cartilage and bone instead of meat in heated meat products is considered as fraud. Due to the economic value of meat, the likelihood of using this unauthorized tissue is possible and

1. Examine the marketed chicken luncheon sausage for its quality and safety for human consumption through assessment of bacteriological quality with special reference to food poisoning microorganisms.

2. Analyze the sensory parameters and deterioration criteria to assure quality in the

1. Materials and methods

Market survey

A total of ninety Egyptian chicken luncheon sausage produced by six different meat processing plants (Fifteen samples each) were collected from different production lots within one month after processing from the supermarkets in Giza and Cairo. Samples were transferred to the laboratories of Food Hygiene

Investigations

1. Sensory panel analysis

Sensory panel analysis was performed by 5 panelists from the members of Food Hygiene and Control Department, Faculty of Veterinary medicine, Cairo University. All chicken luncheon sausage samples were evaluated for appearance, flavor, taste, consistency and

2. Deterioration criteria

2.1. pH value

Five grams from each of the prepared sample were homogenized with 20 ml distilled water for 10-15 seconds (Kandeepan et al., 2009). The pH was measured using pH meter

2.2. Thiobarbituric Acid value

Five grams from each sample were homogenized with 15 ml deionized distilled water. One milliliter of the homogenate was mixed with 50 µl butylated hydroxyanisole (7.2%) and one ml each of 15mM 2-thiobarbituric acid and 15% trichloroacetic acid. The mixture was vortexed, incubated in a boiling water bath for 15 minutes to develop

2.3. Total Volatile Basic Nitrogen

Ten grams sample were macerated with 100 ml tap water and washed into a distilling flask with 200 ml tap water, then 2 grams magnesium oxide were added. A macro-Kjeldahl distillation apparatus was connected

formulation used in the preparation of meat products does not respect the standard and hygiene food regulation. Moreover, addition of inedible offals such as udders, lung, spleen, etc into most products which is sometimes used by producers only can be detected histologically (Rezaian and Rokni, 2003 and Latorre et al., 2015). Therefore, the present study was designed to:

aspects of consumer acceptability and degree of freshness.

3. Detect the presence of bone and cartilage of MDM and the non meat tissues through histological examination.

4. Compare the obtain results with the Egyptian standard specification of chicken luncheon.

and Control Department, Faculty of Veterinary Medicine, Cairo University and kept chilled at 4°C till the time of investigation. All samples were subjected to sensory investigation, determination of deterioration criteria, bacteriological examination as well as detection of non-meat tissues by histological examination.

overall acceptability using a 7-point scale (where 7 denote extremely acceptable and 1 denotes extremely unacceptable) according to the method described by AMSA (1995). Prior to the analysis panelists were trained in the definition and intensities of all investigated sensory parameters.

(Lovibond Senso Direct) with a probe type electrode (Senso Direct Type 330) where three reading for each sample were obtained and the average was calculated. The pH meter was calibrated every two samples using two buffers pH 7.0 and 4.0.

color, then cooled and centrifuged for 15 min at 2500 rpm. The absorbance of the resulting supernatant was measured at 531 nm using Unico 1200 (USA) series spectrophotometer against a blank containing 1 ml of deionized water and 2 ml of 2-thiobarbituric acid-trichloroacetic acid solution. TBA (mg/malonaldehyde/Kg) reading was multiplied by 7.8 (Du and Ahn, 2002).

to the distillation flask containing 25 ml of 2% boric acid solution and few drops of methyl-red indicator (0.016 g methyl red, 0.083 g bromocresol green per 100 ethanol) with the receiving tube was dipped below the liquid, with distillation continued till collection of 200

ml. The condenser was then washed with distilled water and the distillate was titrated with 0.05 M (0.1N) sulphuric acid. The Total

3. Bacterial examination

The spreading technique and the standard plate count agar were used for enumeration of aerobic mesophilic count (Swanson et al., 1992), and aerobic psychrotrophic count (Cousin et al., 1992). While, reinforced clostridial agar and the anaerobic system were performed for enumeration of the anaerobic sporeformers (Brewer and Allgeier, 1966). Coliforms were enumerated "MPN" using lauryl sulfate tryptose broth according

4. Histological examination

A duplicate 1x1 cm blocks from all the investigated samples were fixed in 10% formal saline for twenty four hours followed by overnight washing with running water. Fixed samples were dehydrated in a chain of

Statistical analysis

All data were analyzed using SPSS statistics 17.0 for windows and results were recorded as mean±SE. Analysis of variance was performed by ANOVA procedure to compare between

2. Results and discussion

Table (1): Sensory panel scores of Egyptian chicken luncheon sausage produced by different processing plants (n=90):

	Appearance	Flavor	Taste	Consistency	Overall acceptability
	Mean±SE				
1	2.81 ^{ac} ±0.29	2.64 ^{ac} ±.30	2.38 ^{ac} ±0.27	2.92 ^{ad} ±0.29	2.69 ^a ±0.22
2	2.09 ^a ±0.19	1.82 ^a ±0.26	1.73 ^a ±0.16	2.05 ^b ±0.18	1.93 ^b ±0.11
3	3.57 ^{bc} ±0.25	4.24 ^b ±0.31	4.01 ^b ±0.29	3.56 ^{ac} ±0.26	3.90 ^c ±0.24
4	4.29 ^b ±0.42	4.25 ^b ±0.40	4.04 ^b ±0.43	4.00 ^c ±0.43	4.15 ^c ±0.40
5	2.97 ^c ±0.29	2.90 ^c ±0.38	2.41 ^{ac} ±0.24	3.13 ^a ±0.34	2.90 ^a ±0.27
6	2.81 ^{ac} ±0.12	3.11 ^c ±0.31	2.90 ^c ±0.22	2.27 ^{bd} ±0.21	2.82 ^b ±0.12
Mean	3.09±0.13	3.16±0.16	2.91±0.14	2.99±0.14	3.06±0.13

* a-d: Means with different superscripts differ significantly at P<0.05. 1-6: number of processing plants

Table (2): Deterioration criteria of Egyptian chicken luncheon sausage produced by different processing plants (n=90):

	PH	TVBN	TBA
	Mean ±SE		
1	6.71 ^a ±0.05	5.97 ^a ±0.52	1.11 ^a ±0.08
2	6.51 ^{ab} ±0.04	5.90 ^a ±0.49	1.14 ^a ±0.10
3	6.43 ^b ±0.09	8.14 ^b ±0.43	0.94 ^{ab} ±0.11
4	6.49 ^b ±0.09	6.33 ^{ac} ±0.57	0.86 ^b ±0.03
5	6.36 ^{bc} ±0.07	7.39 ^b ±0.55	1.00 ^{ab} ±0.07
6	6.19 ^c ±0.11	5.41 ^a ±0.33	1.09 ^{ab} ±0.10
Mean	6.45±0.03	6.52±0.22	1.02±0.04

Volatile Base Nitrogen (mg/100 gram sample) was calculated as the titration multiply by 14 (Kearsley et al., 1983).

to Hitchins et al. (1992), faecal coliforms "MPN" were calculated according to FAO (1992) and E. coli was isolated by inoculation on EMB according to Krieg and Holt (1984). For Salmonella isolation, the technique adopted by FAO (1992) was used by pre-enrichment in buffered peptone water 1% followed by selective enrichment in Rappaport medium and selective plating on both S.S. and XLD agar.

upgrading concentration of ethyl alcohol, cleaned in xylene, and embedded in paraffin. Paraffin blocks were sectioned at 4-6 um thickness, and stained with Haematoxyline and Eosine (Banchroft et al., 1996).

chicken luncheon sausages of different processing plants by the least significant difference (LSD) and significance was defined at P<0.05.

* a-c: Means with different superscripts differ significantly at P<0.05.1-6: number of processing plants TVBN expressed as mg/100g sample. TBA expressed as milligrams of malonaldehyde/kg

Table (3): Bacterial load (log₁₀ CFU/g) of Egyptian chicken luncheon sausage produced by different processing plants (n=90):

	Aerobic plate count	Psychrotrophes	Anaerobes
	Mean ±SE		
1	5.31 ^{a,c} ±0.14	5.44 ^a ±0.18	3.73 ^a ±0.14
2	5.79 ^a ±0.18	5.68 ^a ±0.15	3.43 ^{a,c} ±0.13
3	5.15 ^{c,b} ±0.21	5.48 ^a ±0.16	3.66 ^a ±0.20
4	4.73 ^{b,d} ±0.21	4.50 ^b ±0.14	3.00 ^{b,c} ±0.20
5	5.19 ^{c,d} ±0.19	4.77 ^b ±0.21	2.81 ^b ±0.12
6	4.69 ^b ±0.11	4.88 ^b ±0.18	2.79 ^b ±0.22
Mean	5.14±0.08	5.12±0.08	3.24±0.08

* a-c: Means with different superscripts differ significantly at P<0.05, coliforms, fecal coliforms, salmonella and E.coli were not detected in any sample.1-6: number of processing plants




Photo (1): Histological sections stained with H&E (x40) of market Egyptian chicken luncheon sausage. First (A-B); Second (C-D); Third (E-F); Forth (G-H); Fifth (I-J) and Sixth meat processing plant (K-L). Feather (Fe); Skin (S); Fat (F); Blood vessel (Bv); Muscle (Ms); digestive tract (D); Cartilage (C); Tendon (T); Glandular structure (Gl) and Bone (B).

considered to be the most important determinant factor of consumer acceptability. Moreover, sensory analysis permits assessing sensory properties which are directly related with quality perception by consumers (Skinner and Rao, 1986). Data in Table 1 summarized results of sensory evaluation of chicken luncheon sausage collected from six different meat processing plants. It was evident that the overall acceptability of all examined samples was obviously low

Product quality and safety is probably the most important aspect of making poultry meat products because it addresses the question of consumer acceptance and public health safety. While a processor may produce a wonderful sausage, the product must ultimately satisfy the consumer in terms of appearance, color, texture, and flavor (Toldrá et al., 2007 and Jayasena et al., 2013). In this concern AMSA (1984) pointed out that sensory attributes of meat and meat products are widely

higher overall acceptability score but still unacceptable (3.90 and 4.15 respectively).

may not meet its expected sensory, microbiological and nutritional attributes. Furthermore, the unacceptable flavor and lower degrees of acceptability may be due to oxidation of the higher polyunsaturated fatty acids in chicken meat (Jayasena et al., 2013 and Amensour et al., 2015) or due to the use of low quality meat in the processing of such product (Sallam et al., 2004). At the same time, AL-Dughaym and Altabari, (2010) found that thirty percentages of chicken frankfurter not comply with Saudi Standards due to sensory unacceptability. While Gab-Allah (1990) reported that 100% of chicken luncheon were organoleptically acceptable.

the maximum limit for (0.9mg malonaldehyde/kg sample) according to ESS (1696/2005), where the samples had values ranged from 0.086 to 1.14 with an average of 1.02 (mg malonaldehyde/kg). This may be due to fat oxidation or due to the use of low quality meat ingredients in the processing of such products and/or to the high level of lipids and unsaturated fatty acids present. Formulation of luncheon sausage with MDM may promote lipid oxidation of the product (Ertas, 1998), because MDM contains more polyunsaturated fatty acids and haemoproteins which are more susceptible to both chemical and biochemical oxidation. Moreover, heat generation during production of MDM enhances fat rancidity (Püssa et al., 2008). Meanwhile, Andreo et al. (2003) found that thermal process of meat emulsion tend to promote lipid oxidation by disrupting cell membranes and releasing pro-oxidants, thereby inducing the development of 'warmed-over' flavor. The present result was in disagreement with Al-Abdullah and Al-Majali (2011) who concluded that the oxidative rancidity of all examined chicken meat luncheon samples were less than the threshold level.

5.79 with an average of 5.14 (CFU/g). While, psychrotrophic bacterial counts of chicken luncheon sausage were high in all examined processing plants where the first, second and third processing plant showed significant higher values than other processing plants. Moreover, anaerobic bacterial counts ranged

decrease in all the investigated sensory attributes. The product of the third and fourth processing plant scored significantly ($P < 0.05$) The obtained results were in agreement with Sitz et al. (2005) who reported that the overall acceptance of meat products depends on their flavor which is mainly determined by taste and odor compounds. Unacceptable flavor was predominant in most examined samples, which may be explained on the basis of addition of high amount of non meat tissue such as mechanically deboned meat (Photo 1). This non meat tissue is liable for rapid onset of oxidative rancidity resulted in off-flavors and off-odors that ultimately reduced the consumer acceptability (Field, 1988). Moreover, Hargin (1996) and Lumley (1996) stated that undeclared additions of non meat tissues at the expense of skeletal muscle to meat products

Concerning the deterioration criteria of chicken luncheon sausage samples (Table 2), the relatively higher alkaline pH was observed in most processing plant with a mean value (6.45). This may be attributed to metabolites accumulation through bacterial action of spoilage microorganism on protein and amino acids (Kumar et al., 2011). Moreover, Pereira et al. (2011) attributed the higher pH value of emulsion type sausage to higher MDM content. El-Khateib et al. (1988) reported higher pH values (6.3) and El-deeb et al. (2011) found lower pH values (6.08) for chicken luncheon. The results of total volatile base nitrogen were ranged from 5.41 to 8.14 with an average of 6.52 (mg/100g). The higher TVBN value was recorded in the third (8.14) and fifth processing plant (7.39). Low TVBN of examined samples may refer to its low protein content (Farouk et al., 2002). As most of meat processors use MDM as an inexpensive material to substitute meat either partially or even totally in formulation of poultry and meat products (Emara, 2005). The obtained results were confirmed with histological examination where all photo filled with bone, cartilage and skin. Most of TBA values of the examined samples were exceeded Data in Table 3 summarized results of bacterial load (\log_{10} CFU/g) of Egyptian chicken luncheon sausage samples. Total aerobic mesophilic bacterial counts were exceeded the limit established by ESS (1696/2005) (not more than 10^4 CFU/g). Where the samples had values ranged from 4.69 to

bacterial counts in chicken luncheon sausage samples might be attributed to incorporation of high amount of non meat tissue such as MDM, skin with feather and digestive tract (Photo 1).

and handled) and press for elimination of pathogens before the product reaches the consumer. This creates a challenge to the poultry industry to improve the microbiological safety and quality of its products (Shackelford et al., 1988). The present results for APC nearly similar to those obtained by Essa et al. (2004); El-deeb et al. (2011) and Ibrahim et al. (2014). While, lower results were recorded by Sharaf and Sabra (2012) and higher results were recorded by Zahran et al. (2008); AL-Dughaym and Altabari (2010); Shawish (2011) and Nabil et al. (2014). Moreover, lower result than the present result of psychotropic and anaerobic count was recorded by Bkheet et al. (2007) and Nabil et al. (2014) in chicken luncheon, respectively.

processing which injured these sensitive bacteria, such as heat treatment or use of antimicrobial substances as chlorine components and sorbates. However, Luiz et al. (2004) confirmed that Salmonella was able to survive in chicken meat emulsion after the addition of preservatives a fact that could signify a serious risk in the final product and failed to isolate from chicken frankfurter sausages due to adequate industrial cooking process. Similar findings regarding isolation of E.coli were reported by Edris et al. (1992); Ebeid (1996); El-Sbagh (2010) and Shawish (2011). On the other hand, Sharaf and Sabra (2012) isolated E. coli from chicken luncheon with incidence 25%. The present result concerning isolation of Salmonellae was in agreement with AL-Dughaym and Altabari (2010); Shawish (2011); and Ibrahim et al. (2014). However, Essa et al. (2004) isolated 2 strains of Salmonella serotyped as S. Typhimurium and S. Typhi from chicken luncheon.

and Nouman, 2002 and Tremlova and Štarha, 2003). Therefore, it is suitable method for the meat quality control (Buche and Mauron, 1997). The results of histological examination (Photo 1) showed that low skeletal muscular tissues and high fat content were evident in all

from 2.81 to 3.73 with an average of 3.24 CFU/g. ESS (1696/2005) for Egyptian chicken luncheon sausage doesn't include anaerobic bacterial count as quality indicator parameter. The marked increase of all investigated Several authors explained the high bacterial loads to mishandling, incorporation of contaminated raw material, contact with insanitary equipment and insufficient heat treatment during processing as well as unsuitable condition during storage (Cox et al., 1998; Zahran et al., 2008 and Shawish, 2011). In addition to, poultry products offer ideal medium for microbial growth due to their suitable chemical composition and favorable pH a matter which suggested that an improvement of the microbiological quality of poultry meat is necessary. Luiz et al. (2004) recorded a considerable number of spoiled chicken sausages delivered to retailers before the recommended sell-by date incurring large financial losses. Therefore, when contamination and illness occurred, investigators tend to look at the product (how it was produced, processed Regarding our results of coliforms counts, it is failed to be detected in all examined chicken luncheon sausage samples. However, El-deeb et al. (2011) recorded low contamination rate with coliforms in chicken luncheon and frankfurter (3.4×10 and 3.9×10 respectively Cfu/g). Zhang et al. (2001) explained that to good packaging, good retailing and no handling by workers, in addition to their exposure to heat treatment and smoking during processing. On other hand, Bkheet et al. (2007); Shawish (2011) and Nabil et al. (2014) observed high contamination rate of coliforms in chicken luncheon. Moreover, both E. coli and Salmonella failed to be isolated from all examined samples and it may be due to sufficient cooking temperature of such product. According to ESS (1696/2005) for Egyptian chicken luncheon sausage it must be free from both E. coli and Salmonella. Morrison and Fleet (1985) attributed the absence of Salmonellae in chicken meat products to the different Histological examination seems to be very accurate and rapid technique (lasts one day only) that can be used effectively to detect adulteration of meat products such as Egyptian luncheon as an emulsion type product with avian skin even in products subjected to severe treatments (Emara

salivary gland tissue, nuchal ligament (Rokni et al., 1997), chicken skin, hyaline cartilage, peritoneal fat and kidney (Sepehri, 2008) in heated sausage. Moreover, Latorre et al. (2015) concluded that the formulations used in the preparation of the examined products do not respect the standard and hygiene food regulation and the products are of bad quality.

histological examination confirmed the use of unauthorized tissue in formulation of such product which does not respect the standard and hygiene food regulation. Therefore, based on this evidence, control of the formulation process along with detailed monitoring and continuous inspection of this product manufacture is essential to achieve its quality and safety.

AMSA "American Meat Science Association" (1984): American Science Association Guidelines for Sensory, Physical and Chemical Measurements of Ham. National Livestock and Meat Board, Chicago, USA.

AMSA "American Meat Science Association" (1995): Research guidelines for cookery, sensory evaluation and instrumental tenderness measurements of fresh meat. Chicago, IL, USA:.

Andreo, A.I.; Doval, M.M.; Romero, A.M and Judis, M.A. (2003): Influence of heating time and oxygen availability on lipid oxidation in meat emulsions. *European Journal of lipid science and Technology*, 105(5): 207-213.

Banchroft, J.D.; Stevens, A. and Turner, D.R. (1996): Theory and Practice of Histological Techniques, 4th Ed. Churchill Livingstone, New York, London, San Francisco, Tokyo.

Barbut, S. (2002): Poultry Products Processing. An Industry Guide, 2nd Ed. CRC Press. Boca Raton, London, New York, Washington, D.C.

Bkheet, A.A.; Rezk, M.S.H. and Mousa, M.M. (2007): Study on the microbiological content of local manufactured poultry meat products in

samples. Moreover, bone particles, cartilage particles, skin tissue and other non meat tissue such as digestive tract were observed. All these non meat tissues have negative impact on the quality and safety of the examined product which confirmed the results of sensory (Table 1), deterioration (Table 2) and bacteriological investigation (Table 3). Several authors observed presence of non meat tissue like

4. Conclusions

This study has shown that most of the investigated chicken luncheon sausages were not copy with the Egyptian Standard Specifications for chicken luncheon in term of sensory attributes, deterioration criteria as well as very high bacteriological load. Chicken luncheon sausages samples were highly contaminated with aerobic and anaerobic bacterial counts while *E. coli* and *Salmonella* failed to be isolated. Moreover, the results of

References

- Agamy, N.F. and Hegazy, E.M. (2011):** Quality and safety evaluation of marketed breaded chicken and production of a high quality Nuggets product. *Australian Journal of Basic and Applied Science*, 5(9): 661-671.
- Al-Abdullah, B. and Al-Majali, A. (2011):** Effects of fat content and heat treatment on the chemical and sensory characteristics of canned luncheon meat. *Jordan Journal of Agricultural Sciences*, 7(4): 701-709.
- AL-Dughaym, A.M. and Altabari, G.F. (2010):** Safety and quality of some chicken meat products in Al-Ahsa markets-Saudi Arabia. *Saudi Journal of Biological Sciences*, 17(1): 37-42.
- Amensour, M.; Sendra, E.; Pérez-Alvarez, J.Á.; Abrini, J. and Fernández-López, J. (2015):** Effect of myrtle (*myrtus communis*) extracts on storage stability of chicken frankfurters. *International Journal of Biotechnology for Wellness Industries*, 4(1): 1-11.
- American Meat Institute (2004):** Overview of U.S. Meat and Poultry Production and Consumption. <http://www.meatami.com/>

- Veterinary Medicine, Zagazig University, (Benha Branch).
- Edris, M.A.; Abd El-Rahman, M. and Mousa, M. M. (1992):** Microbial evaluation of some retailed cut-up chicken and poultry meat products. *Benha Veterinary Medicine Journal*, 3 (1): 154-166.
- El-deeb, M.F.; El-Glel, H.A. and Samaha, I.A.T. (2011):** Quality assurance of some poultry meat products. *Alexandria Journal of Veterinary Science*, 33(1): 153-163.
- EL-Khateib, T.; Abd El-Rahman, H.; Hamdy, M. and Lotfi, A. (1988):** Poultry meat products in Egypt "Proximal chemical composition and microbiological quality". *Fleischwirtschaft*, 68 (6): 756-757.
- El-Sbagh, R.A.E. (2010):** Escherichia coli in meat and poultry products with special reference to identification of Verotoxigenic Escherichia coli using the PCR technique. M.V.Sc., Thesis, Faculty of Veterinary Medicine, Benha University.
- Emara, M.M.T. (2005):** Use of chlorine, benzoate and lactic acid as decontaminators to mechanically separated poultry meat and application in traditional Egyptian luncheon. *Veterinary Medical Journal, Giza*, 53(1): 153-162.
- Emara, M.M.T. and Nouman, T.M. (2002):** Histological detection of avian skin in meat products and its impact on quality. *Egyptian Veterinary Medical Association*, 62(1): 147-156.
- Ertaş, A.H. (1998):** The oxidation of meat fats. *Gıda*, 23 (1): 11-17.
- ESS 1696 (2005):** chicken luncheon. Egyptian Organization for Standardization and Quality Control.
- Essa, H.H.; Maker, N.H.; Sohair, A. and Hussein, Z. (2004):** Bacteriological evaluation of chicken luncheon in Assiut City. *Assiut-veterinary-Medical-Journal*, 50(102): 64-71.
- FAO "Food and Agricultural Organization" (1992):** Manual of Food and Quality Control. 4: Microbiological Analysis. Food and Agricultural Organization of the United Nation, Rome.
- El-Bohira governorate. Assiut Veterinary Medical Journal, 53(112): 115-125.**
- Bonoli, M.; Caboni, M.F.; Rodriguez-Estrada, M.T. and Lercker, G. (2007):** Effect of feeding fat sources on the quality and composition of lipids of precooked ready-to-eat fried chicken patties. *Food Chemistry*, 101(4): 1327-1337.
- Brewer, J.H. and Allgeier, D.L. (1966):** Self contained carbon dioxide hydrogen anaerobic system. *Applied Microbiology*, 14(6): 985-988.
- Buche, P. and Mauron, D. (1997):** Quantitative characterization of muscle fiber by image analysis. *Computers and Electronics in Agriculture Journal*, 16(3): 189-217.
- Cousin, M.A.; Jay, J.M. and Vasavada, P. (1992):** Psychrotrophic Microorganisms. In: "compendium of Methods for the Microbiological Examination of Foods". Chapter 9, 3rd edition, Pp. 153-164, C. Vanderzant and D.F. Splittstoesser, (Eds). American Public Health Association, Washington, DC.
- Cox, N.A.; Russell, S.M. and Bailey, J.S. (1998):** The microbiology of stored poultry. In: "Microbiology of Meat and Poultry". Pp: 266-87. Davies, A and R. Board (Eds). Blackie Academic and Professional, London.
- Daros, F.G.; Masson, M.L. and Amico, S.C. (2005):** The influence of the addition of mechanically deboned poultry meat on the rheological properties of sausage. *Food engineering*, 68(2): 185-189.
- Deumier, F. and Collignan, A. (2003):** The effects of sodium lactate and starter cultures on pH, lactic acid bacteria, *Listeria monocytogenes* and *Salmonella* spp. levels in pure chicken dry fermented sausage. *Meat Science*, 65(3): 1165-1174.
- Du, M. and Ahn, D.U. (2002):** Effect of antioxidants on the quality of irradiated sausages prepared with turkey thigh meat. *Poultry Science*, 81(8): 1251-1256.
- Ebeid, S.R.B. (1996):** Incidence of some Enterobacteriaceae organism in retailed unpacked raw cut-up chicken. M.V.Sc., Thesis. Meat hygiene, Faculty of

- organoleptic qualities. Tamil nadu Journal of Veterinary and Animal Sciences, 8 (2): 60-67.
- Jayasena, D.D.; Ahn, D.U.; Nam, K.C. and Jo, C. (2013):** Flavour chemistry of chicken meat: A Review. Asian Australas. Journal of Animal Science, 26(5): 732-742.
- Kandeepan, G.; Anjaneyulu, A.S.R.; Kondaiah, N.; Mendiratta, S.K. and Lakshmanan, V. (2009):** Effect of age and gender on the processing characteristics of buffalo meat. Meat Science, 83(1): 10-14.
- Kearsley, M.W.; El-Khatib, L. and Gunu, C.O.K.A. (1983):** Rapid determination of total volatile nitrogen in fish and meat. Association of Public Analysts, 21(4): 123-128.
- Krieg, N.R. and Holt, J.G. (1984):** Bergey's Manual of Systematic Bacteriology. Vol. 1, Williams and Willkins Baltimore, USA.
- Kumar, D. and Tanwar, V.K. (2011):** Effects of incorporation of ground mustard on quality attributes of chicken nuggets. Journal of Food Science and Technology, 48(6): 759-762.
- Latorre, R.; Sadeghinezhad, J.; Hajimohammadi, B.; Izadi, F. and Sheibani, M.T. (2015):** Application of Morphological Method for Detection of Unauthorized Tissues in Processed Meat Products. Journal of Food Quality and Hazards Control, 2(2): 71-74.
- Luiz, A.F.; Moreira, F.C.; Corrêa, E.F. and Falcão, D.P. (2004):** Monitoring of the dissemination of Salmonella in the chicken frankfurt-sausage production line of a sausage factory in the state of são paulo, Brazil. Memórias do Instituto Oswaldo Cruz Rio de Janeiro, 99(5): 477-480.
- Lumley, I.D. (1996):** Authenticity of meat and meat products. In "Food Authentication" Pp. 108-139, P.R. Ashurst and M.J. Denis (Eds). Blackie Academic and Professional, London.
- Morrison, G.J. and Fleet, G.H. (1985):** Reduction of Salmonella on chicken carcasses by immersion treatments. Journal of Food Protection, 48(11): 939-943.
- Nabil, M.E.; Edris, A.M.; Shaltout, F.A. and Zakaria, I.M. (2014):** Effect of Farouk, M.M.; Wielezko, K.; Lim, R.; Turnwald, S. and Macdonal, G.A. (2002): Cooked sausage batter cohesiveness as affected by sarcoplasmic proteins. Meat Science, 61(1): 85-90.
- Field, R.A. (1988):** Mechanically Separated Meat, Poultry and Fish. In "Edible Meat By-Products". Pp. 83-128. A.M. Pearson and T.R. Dutson, (Eds). New York: Elsevier Applied Science.
- Freitas, M.Q.; Silva, T.J.P.; Mano, S.B. and Chaves, J.B.P. (2004):** Medidas instrumentais de textura e cor, em mortadela produzida com carne mecanicamente separada de frango. Higiene Alimentar, 18(126/127), 66-70.
- Gab-Allah, H.M.H. (1990):** Sanitary status of some meat poultry products marketed in sharkia Governorate. M.V.Sc., Thesis in Meat Hygiene, Faculty of Veterinay Medicine, Zagazig University.
- Gill, C.O. (1988):** Microbiology of edible meat by products. In "Edible Meat By-products". Pp. 47-82. A.M. Pearson and T.R. Dutson (Eds). Barking, Elsevier Applied Science.
- Guerrero-Legarreta, I. and Hui, Y.H. (2010):** Processed poultry products: A primer. In "Handbook of Poultry Science and Technology". Volum 2, Pp. 3-10, Wiley.
- Hargin, K.D. (1996):** Authenticity issues in meat and meat products. Meat Science, 43(13): 277- 289
- Hitchins, A.D.; Hartman, P.A. and Todd, E.C. (1992):** Bergey's Manual of Systematic Bacteriology. Vol. 1, Williams and Willkins Baltimore, USA.
- Hwang, K.E.; Choi, Y.S.; Choi, J.H.; Kim, H.Y.; Kim, H.W.; Lee, M.A.; Chung, H.K. and Kim, C.J. (2011):** Effect of ganghwayakssuk (*Artemisia princeps* Pamp.) on oxidative stability of deep fried chicken nuggets. Food Science Biotechnology, 20(5): 1381-1388.
- Ibrahim, H.M.; Amin, R.A.; Ibrahem, I.A. and Yunis, O.F. (2014):** Isolation of Enterobacteriaceae from poultry products in El-Behera and Alexandria governorates. Benha Veterinary Medical Journal, 27(1): 109-117.
- Ilayabharathi, D., Sheriff, F.R. and Manohar, G.R. (2012):** Shelf-life of spent chicken sausage and its

- sausage. *Lebensmittel-Wissenschaft und-Technologie*, 37(8): 849-855.
- Sepehri, E.S. (2008):** Histological methods evaluation for detection of adulteration of raw meat products supplied in Tehran. Ph.D., Thesis, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran.
- Shackelford, A.D. (1988):** Modification of processing methods to control *Salmonella* in poultry, *Poultry Science*, 67(6): 933-935.
- Sharaf, E.M. and Sabra, S.M. (2012):** Microbiological loads for some types of cooked chicken meat products at Al-Taif Governorate, KSA. *World Applied Sciences Journal*, 17 (5): 593-597.
- Shawish, R.R.M. (2011):** Microbial evaluation of some retailed cut-up chicken and poultry meat products. M.V.Sc. Thesis, Faculty of Veterinary Medicine, Sadat city branch, Menoufia University.
- Sitz, B.M.; Calkins, C.R.; Feuz, D.M.; Umberger, W.J. and Eskridge, K.M. (2005):** Consumer sensory acceptance and value of domestic, Canadian, and Australian grass-fed beef steaks. *Journal of Animal Science*, 83(12): 2863-2868.
- Skinner, G.E. and Rao, V.N.M. (1986):** Linear viscoelastic behavior of frankfurters. *Journal Texture Studies*, 17(4): 421-432.
- Swanson, K.M.J.; Busta, F.F.; Peterson, E.H. and Johnson, M.G. (1992):** Colony Count Methods., In "Compendium of Methods for The Microbiological Examination of Foods". Chapter 4, Pp. 75-77. 3rd edition, C. Vanderzant and D.F. Splittstoesser, (Eds). American Public Health Association, Washington, DC.
- Toldrá, F.; Hui, Y.H.; Astiasarán, I.; Nip, W.; Sebranek, J.G.; Silveira, E.F.; Stahnke, L.H. and Talon, R. (2007):** Handbook of Fermented Meat and Poultry. First edition, Blackwell Publishing, USA.
- Tremlova, B. and Starha, P. (2003):** Histometric evaluation of meat products - determination of area and comparison of results obtained by histology and chemistry. *Czech Journal Food Science*, 21(3): 101-106.
- some preservatives on bacterial load of some poultry meat products. *Benha Veterinary Medical Journal*, 26(1): 94-103.
- Park, K.S.; Choi, Y.S.; Kim, H.Y.; Kim, H.W.; Song, D.H.; Hwang, K.E.; Choi, S.G. and Kim, C.J. (2012):** Quality characteristics of chicken emulsion sausages with different levels of makgeolli lees fiber. *Korean Journal of Food Science Animal Resource*, 32(1): 54-61.
- Pereira, A.G.T.; Ramos, E.M.; Teixeira, J.T.; Cardoso, G.P.; Ramos, A.L.S. and Fontes, P.R. (2011):** Effects of the addition of mechanically deboned poultry meat and collagen fibers on quality characteristics of frankfurter-type sausages. *Meat Science*, 89(4): 519-525.
- Potter, N. (2001):** Food Science. 4th Ed., The AVI publishing Co., Inc., New York, USA.
- Püssa, T.; Pällin, R.; Raudsepp, P.; Soidla, R. and Rei, M. (2008):** Inhibition of lipid oxidation and dynamics of polyphenol content in mechanically deboned meat supplemented with sea buckthorn (*Hippophae rhamnoides*) berry residues. *Food Chemistry*, 107(2): 714-721.
- Püssa, T.; Raudsepp, P.; Toomik, P.; Pällin, R.; Mäeorg, U.; Kuusik, S.; Soidla, R. and Rei, M. (2009):** A study of oxidation products of free polyunsaturated fatty acids in mechanically deboned meat. *Food Composition and Analysis*, 22(4): 307-314.
- Rezaian, M. and Rokni, N. (2003):** Histological study of the heated meat products of mazandaran province of Iran. The 11th international symposium of the world association of veterinary laboratory diagnosticians and OIE seminar on biotechnology
- Rokni, N.; Rezaian, M. and Dayani D.A. (1997):** Histological and histometrical study of different heated sausages. *Journal of Veterinary Medicine, Tehran University*, 52(2): 95-103.
- Sallam, Kh.I.; Ishioroshi, M. and Samejima, K. (2004):** Antioxidant and antimicrobial effects of garlic in chicken

Zhang, L.; Davis, M.A. and Conner, D.E. (2001): Poultry-borne pathogens: plant considerations. In "Poultry Meat processing" Chapter 9. A.R. Sams (Ed). CRC Press LLC, New York, USA.

Zahran, D.A.; Hendy, B.A. and El-Hifnawi, H.N. (2008): Incidence and radiation sensitivity of bacillus cereus, listeria monocytogenes and their toxins in some chicken products. World Applied Science Journal, 5(2): 182-188.

الملخص العربي

تم جمع عدد تسعين عينة من لانشون لحم الدجاج المنتج محالياً من ستة شركات مصنعات اللحوم المختلفة بواقع خمسة عشر عينة من كل شركة. فقد تم جمعها بصورة عشوائية في خلال الشهر الأول بعد تصنيعها وفحصها لتحديد مدى جودتها. وأجريت على هذه العينات اختبارات الفحص الحسي، وفحوصات دلالات الفساد، الفحص الميكروبيولوجي بالإضافة إلى الكشف على وجود مخلفات أخرى خلاف لحم الدجاج بالفحص المجهرى باستخدام الميكروسكوب الضوئي. فقد دلت نتائج فحص تلك العينات على تدهور الخواص الحسية عن الصورة المتعارف عليها لمصنعات لانشون الدواجن بدرجة كبيرة بالإضافة إلى وجود رائحة التزنخ الطاغية على المنتج. بينما دلت فحوصات دلالات الفساد على ارتفاع درجة الأس الهيدروجيني وحامض الثيوباربيتورك وهذا يؤكد نتائج الفحص الحسي. كما أشارت نتائج الفحص الميكروبيولوجي إلى زيادة العد الكلي للبكتريا الهوائية واللاهوائية بينما لم يتم عزل كلا من الايشيريشياكولاى والسالمونيلا. وقد أظهر الفحص المجهرى باستخدام الميكروسكوب الضوئي ان معظم العينات التى تم فحصها تحتوي على نسبة هائلة من النسيج الضام، العظام، الغضاريف، الجلد مما يؤثر سلباً على الجودة الحسية وكذلك المحتوي الميكروبي لهذا المنتج. وقد لخصت هذه الدراسة على ان معظم مصنعات لانشون الدجاج لا تطابق المواصفه القياسيه المصريه.