

SEROLOGICAL DIAGNOSIS OF *NEOSPORA CANINUM* INFECTION IN SOME DOMESTIC ANIMALS FROM EGYPT

A. EL-GHAYSH*, FATHIA KHALIL**, M. HILALI* and A. M. NASSAR

*Parasitology Department, Faculty of Veterinary Medicine, Cairo University, Egypt

** National Research Center, Dokki, Egypt

Received: 1.3.2003.

Accepted: 4.5.2003.

SUMMARY

This study investigates the presence of antibodies to *N. caninum* in cattle, sheep, goats and dogs from Egypt. Sera from the tested animals were examined using a direct agglutination test incorporating mercaptoethanol and formalin-preserved whole *N. caninum* tachyzoites as antigen. Anti-*N. caninum* antibodies were found in 18 out of 111 aborted cattle (16.2 %). Abortion occurs during the third to ninth month of gestation period. Three of the 26 non-aborted cattle (11.5 %) were seropositive. Statistical analysis indicated that anti-*N. caninum* antibodies occurred more in the aborted than in the non-aborted group (slightly significant). Seventy three of 202 sheep (36.1 %) and 31 of 88 (35.2%) goats recorded antibodies for *N. caninum*. Eight out of 29 (27.6 %) stray dogs had antibodies against *N. caninum*. The present communication is the first nation wide

survey on serological evidence of *N. caninum* antibodies in different life stock and dogs in Egypt.

INTRODUCTION

Neospora caninum is a protozoan parasite that was first recognized in dogs with neurological signs (Dubey et al., 1988). Subsequently the parasite was incriminated as a major cause of abortion in cattle all over the world (Dubey and Lindsay, 1996).

Natural infection with *Neospora species* was reported from sheep, goats and horses (Dubey and Lindsay, 1996). Recently the domestic dog was found to be the definitive host of *N. caninum* since it excretes environmentally resistant oocysts in faeces after ingestion of tissue cysts (Mc Allister et al., 1998). Since accurate diagnosis of *N. caninum* is key for understanding of the

epidemiology of neosporiosis, several reports were published from different countries on the seroprevalence of the parasite in farm animals (Hemphill et al., 2000).

In Egypt there are only two published reports on the seroprevalence of anti-*Neospora caninum* antibodies in camels (Hilali et al., 1998) and water buffaloes (Dubey et al., 1998). Similar studies on other farm animals are non-existent. Therefore, this research was undertaken to estimate the seroprevalence of *N. caninum* antibodies in cattle, sheep, goats and dogs.

MATERIALS AND METHODS

1- Animals and Blood samples

Blood samples were obtained from 111 aborted and 26 non-aborted cattle. Both groups were chosen from 3 dairy Holstein cattle farms (Table 1). All investigated cattle varied in age from 3-7 years old. The gestational age of abortion varied in these farms from 3-9 months. The non-aborted cattle were pregnant at a gestational age varying from 3-9 months. Serum samples were collected immediately after abortion from aborted cattle. While, the samples were collected from the non-aborted cattle during their gestational period.

Two hundred and two sheep were sampled from farms in Delta (183 samples) and in the suburbs of Cairo (19 samples). The age of

sheep varied from 1-3 years. A total of 88 goat blood samples were obtained from farms in the suburbs of Cairo (14), Delta (24) and Sinai Peninsula (50). The goats varied in their ages from 1-3 years.

All the collected dog samples (29) were obtained from stray dogs living in dairy farms in the suburbs of Cairo city. Five ml of blood was obtained from each animal and serum was separated and kept at -20°C until used.

2- Serological analysis

Sera were tested by using *Neospora caninum* agglutination test (NAT) kit (Vetoquinol, France) as described by Romand et al. (1998). The used antigen was formalin-preserved whole tachyzoites of *N. caninum* incorporated with mercaptoethanol. The tested sera in addition to the positive and negative control sera were used at a dilution of 1:40. The samples which showed a diffused precipitation in the bottom of the wells were considered positive while negative samples showed a spot precipitate. The positive samples were serially diluted beginning from 1:40 to 1:640 for the determination of end point titer.

RESULTS

Abortion in cattle with anti-*Neospora* antibodies occurred at a gestational age varying from 3-9 months. As illustrated in table 1, serological examination of 111 aborted cattle showed that only

8 had anti-*N. caninum* antibodies (16.2%). The distribution of these serologically positive cattle was 7 (21.9%), 8 (20%), 3 (7.7%) three different farms. On the other hand, serological examination of pregnant non-aborted cattle chosen from the same farms indicated that 3 cattle (11.5%) from a total number of 26 were positive for *N. caninum*. One of these positive cattle was from farm I and two were from farm II. None of the selected 9 cattle in farm III had antibodies against *N. caninum*. The end point titers among 18 serologically positive aborted cattle were 1:40 (11), 1:80 (2), 1:160 (3) and 1:640 (2) (Table 4). The serologically positive non-aborted cattle had end point titer of 1:40 (two cattle) and 1:80 (one

cow). As shown in table 2 statistical analysis indicated that anti-*N. caninum* antibodies occurred more in the aborted than in the non-aborted group (slightly significance).

Table 3 showed that 73 out of 202 examined sheep (36.1%) had antibodies against *N. caninum*. Sixty seven sheep out of 183 (36.6%) in Delta and 6 of 19 sheep (31.6 %) from Cairo recorded positive results for *N. caninum* antibodies. Five out of the 14 goat samples (35.7%) from Cairo, 8 out of 24 goats (33.3%) in Delta and 18 out of 50 goats (36%) from Sinai reported positive results for *Neospora caninum*. The total percentage of infected goats was 35.2%. Eight

Table 1: Distribution of serologically positive cattle in the 3 farms .

Farm No.	Serological results of aborted cattle					Serological results of pregnant non aborted cattle				
	Total Number	No. of Positive	%	G.A. (month)	No.	No. of examined	No. of Positive	%	G.A. (month)	No.
I	32	7	21.9	5	3	7	1	14.3	4	1
				6	2					
				7	1					
				8	1					
II	40	8	20	4	1	10	2	20	6	1
				5	2				8	1
				6	2					
				7	2					
				8	1					
III	39	3	7.7	5	1	9	Zero	0		
				6	1					
				7	1					
Total	III	18	16.2			26	3	11.5%		

Table 2: chisquare at degree of freedom (3x6) at P> 0.05 of the examined cattle.

GA (month)	Aborted cattle			Non aborted			Total examined
	Positive	Negative	Total	Positive	Negative	Total	
3	0	4	4	0	3	3	7
4	1	12	13	1	4	5	18
5	6	25	31	1	4	5	36
6	5	27	32	0	5	5	37
7	4	16	20	0	2	2	22
8	2	9	11	1	3	4	15
9	0	0	0	0	2	2	2
Total	18	93	111	3	23	26	137

Chisquare all 21.636

Table 3: Results of serological examination of sheep, goats and dogs for *N. caninum*.

Animal	Location	Total Number	Positive	Percentage
Sheep	Delta	183	67	36.6
	Cairo	19	6	31.6
Total sheep		202	73	36.1
Goat	Cairo	14	5	35.7
	Delta	24	8	33.3
	Sinai	50	18	36
Total goat		88	31	35.2
Dog	Cairo	29	8	27.6

Table 4: End point titers for *N. caninum* antibodies in seropositive cattle, sheep, goats and dogs.

	Total	1:40		1:80		1:160		1:320		1:640	
		No	%	No	%	No	%	No	%	No	%
Aborted cattle	18	11	61.11	2	11.11	3	10.66	0	0	2	11.11
Non aborted cattle	3	2	66.66	1	3.8	0	0	0	0	0	0
Sheep	73	61	83.56	10	4.9	0	0	2	2.73	0	0
Goat	31	28	90.32	1	1.3	2	6.45	0	0	0	0
Dog	8	4	50	4	13.8	0	0	0	0	0	0

13 out of 29 dogs (27.6 %) has antibodies against *N. caninum*. The distribution of the end point titers for *N. caninum* antibodies in the seropositive samples is illustrated in Table 4. Among 13 seropositive sheep, 61, 10 and 2 showed end point titer 1:40, 1:80 and 1:320 respectively. While, 28, 1, 2 out of 31 *N. caninum* seropositive goats showed titer 1:40, 1:80 and 1:160 respectively. The seropositive dogs showed end point titer of 1:40 (4) and 1:80 (4).

DISCUSSION

There are several reports on the seroprevalence of antibodies against *N. caninum* in farm animals from different countries by different serological tests (Hemphill et al. 2000). In 1998, two independent papers described *N. caninum* direct agglutination test according to the principle of the *Toxoplasma gondii* modified agglutination test (Romand et al., 1998 and Packham et al., 1998). The test described by Romand et al. (1998) employed tachyzoites of the canine NC-1 isolate while Packman et al. (1998) used a bovine isolate. This study used the bovine isolate (Packham et al., 1998). NAT was proved to be highly specific and sensitive in diagnosis of *N. caninum* in animals infected with related parasites including *T. gondii* (Romand et al., 1998). Because of its simplicity and visibility, the NAT has the potential of being used for analysis of sera from a variety of species, possibly replacing the IFAT (Huong et al., 1998). Hence, in this study we

used NAT as a serological test for detection of *N. caninum* antibodies.

The present study showed that 18 out of 111 (16.2 %) aborted cattle and 3 of the 26 (11.5 %) non aborted cattle had antibodies for *N. caninum*. Statistical analysis indicated that anti-*N. caninum* antibodies occurred more in the aborted than in the non-aborted group (slightly significant). This indicates that *N. caninum* is one of the abortifacient agents in cattle. This agrees with that reported by Anderson et al., (1991) in USA and by Hemphill et al. (2000) in different European countries, New Zealand and Australia.

The incidence of *N. caninum* in cattle was also studied in different countries, Vietnam (5.5%, Huong, et al., 1998); France (5.6%, Ould-Amrouche et al., 1999); Korea (23%, Bae et al., 2000); Sweden (7%, Bjorkman et al., 2000); Brazil (14.09%, Gondim et al., 1999); Thailand (6%, Suteeraparp et al., 1999) and Spain (30.6%, Mainar et al., 1999).

The present study recorded that in Delta and Cairo, 73 of 202 sheep (36.1%) were seropositive for *N. caninum*. Thirty one of 88 (35.2%) goats recorded antibodies for *N. caninum*. Little is known about the seroprevalence of *N. caninum* in sheep and goats. However, Barr et al. (1992) found it in tissue of goats.

Stray dogs showed *N. caninum* antibodies in

7.6%. Cheadle et al. (1999) in Alabama recorded that 11.5% of examined dogs had *N. caninum* antibodies. Wouda et al. (1999) found that 23.6% of dogs in Netherland had been exposed to the parasite. *N. caninum* antibodies were reported from dogs in Australia (9%), Uruguay (20%), Falkland Islands (0.2%), and Belgium (11%) (Barber et al., 1997 a and b). Reichel et al., (1998) found seroprevalence of 22% in dogs from New Zealand, while it was 29% from Italy (Cringoli et al., 1996). The above mentioned variation in seroprevalence rates might be due to the type of the examined dogs (stray or indoor dogs, dogs in dairy farms or in urban areas). The variation between the end point titers in each animal species might be attributed to the difference in age, time of infection and the immune system of the examined animals.

The obtained results from this study could be a preliminary guide for the presence of *N. caninum* in Egyptian animals. Isolation and identification of *N. caninum* from these animals is considered an important step for confirming the role of *N. caninum* in inducing abortion and still birth in Egyptian animals.

REFERENCES

- Anderson, M.L., Blanchard, P.C., Barr, B.C., Dubey, J.P., Hoffman, R.L., Conrad, P.A. (1991). *Neospora*-like protozoan infection as a major cause of abortion in California dairy cattle. *J. Am. Vet. Med. Assoc.* 198, 241-244.
- Barr, B.C., Anderson, M.L., Woods, L.W., Dubey, J.P. and Conrad, P.A. (1992). *Neospora*-like protozoal infections associated with abortion in goats. *J. Vet. Diagn. Invest.* 4, 365-367.
- Barber, J.S., Gasswe, R.B., Ellis, J., Reichel, M.P., McMillan, D., Trees, A.J. (1997a). Prevalence of antibodies to *Neospora caninum* in different canid populations. *J. Parasitol.* 83, 1056-1058.
- Barber, J.S., Van Ham, L., Polis, I., Trees . (1997b). Seroprevalence of antibodies to *Neospora caninum* in Belgian dogs. *J. small Anim. Pract.* 38, 15-16.
- Bae J.S, Kim DY, Hwang WS, Kim JH, Lee NS, Nam HW. (2000). Detection of IgG antibody against *Neospora caninum* in cattle in Korea. *Korean J Parasitol* 38 (4):245-249.
- Bjorkman C, Alenius S, Manuelsson U, UgglA A. (2000). *Neospora caninum* and bovine virus diarrhoea virus infections in Swedish dairy cows in relation to abortion. *Vet J.*, 159(2):201-206.
- Cheadle, M.A., Lindsay, D.S., Rowe, S., Dykstra, C.C., Williams, M.A., Spencer, I.A., Toivio-Kirnnucan, M.A., Lenz, S.D., Newoton, J.C., Rolsoma, M.D. and Blagburn, B.L. (1999). Prevalence of antibodies to *Neospora caninum* in dogs. *Int. J. Parasit.* 29, 1537-1543.
- Cringoli, G., Capuano, F., Veneziano, V., Romano, L., Solimene, R., Barber, J.S. and Trees, A.J. (1996). Prevalence of antibodies against *Neospora caninum* in dog sera. *Parasitologia*, 38, 282.
- Dubey, J.P. and Lindsay, D.S. (1996). A review of *Neospora caninum* and neosporiosis. *Vet. Parasitol.* 67, 1-59.
- Dubey, J.P., Romand, S., Hilali, M., Kwok, O.C.H. and Thulliez, P. (1998). Seroprevalence of antibodies to *Neospora caninum* and *Toxoplasma gondii* in water buffaloes (*Bubalus bubalis*) from Egypt. *Int. J. Parasit.* 28, 527-529.

- Dubey, J. P., Carpenter, J.L., Speer, C.A., Topper, M.J. and Uggla, A. (1988). A Newly recognized fatal protozoan disease of dogs. *J. Am. Vet. Med. Assoc.*, 192, 1269-1285.
- Gondim LF, Sartor IF, Hasegawa M, Yamane I. (1999). Seroprevalence of *Neospora caninum* in dairy cattle in Bahia, Brazil. *Vet Parasitol.* 15;86(1):71-75
- Hemphill, A., Gottstein, B., Conraths, F.J., De Meerschman, F., Ellis, J.T., Innes, E.A., McAllister, M.M., Ortega-Mora, L.M., Tenter, A.M., Trees, A.J., Uggla, A., Williams, D.J.L. and Wouda, W. (2000). A European perspective on *Neospora caninum*. *Int. J. Parasit.*, 30, 877-924.
- Hilali, M., Romand, S., Thulliez, P., Kwok, O.C.H. and Dubey, J. P. (1998). Prevalence of *Neospora caninum* antibodies in sera from camels from Egypt. *Vet. Parasit.*, 75, 269-271.
- Huong, L.T.T., Ljungstrom, B. L., Uggla, A. and Bjorkman, C. (1998). Prevalence of antibodies to *Neospora caninum* and *Toxoplasma gondii* in cattle and water buffaloes in Southern Vietnam. *Vet. Parasit.*, 75, 53-57.
- Mc Allister, M.M., Dubey, J.P., Lindsay, D.S., Jolley, W.R., Wills, R.A. and McGuire, A.M. (1998). Dogs are definitive hosts for *Neospora caninum*. *Vet. Parasitol.* 82, 327-333.
- Mainar-Jaime, R.C., Thurmond, M.C., Berzal-Herranz B. and Hietala, S.K. (1999). Seroprevalence of *Neospora caninum* and abortion in dairy cows in northern Spain. *Vet Rec* 17;145 (3):72-75.
- Ould-Amrouche A, Klein F, Osdoit C, Mohammed HO, Touratier A, Sanaa M, Mialot JP. (1999). Estimation of *Neospora caninum* seroprevalence in dairy cattle from Normandy, France. *Vet Res.* 30(5):531-538.
- Packham, A.E., Sverlow, K.W., Conrad, P.A., Loomis, E.F., Rowe, J.D., Anderson, M.L., Marsh, A.E., Cray, C., Barr, B.C., (1998). A modified agglutination test for *Neospora caninum*: Development, optimization and comparison to the indirect fluorescent-antibody test and Enzyme linked immunosorbent assay. *Clin. Diagn. Lab. Immunol.*, 5, 467-473.
- Reichel, M.P., Thornton, R.N., Morgan, P.L., Mills, R.J.M. and Schar, G. (1998). Neosporosis in a pup. *New Zealand Vet. J.* 46, 106-110.
- Romand, S. Thulliez, P. and Dubey, J.P. (1998). Direct agglutination test for serologic diagnosis of *Neospora caninum* infection. *Parasitol. Res.* 80, 50-53.
- Suteeraparp P, Pholpark S, Pholpark M, Charoenchai A, Chompoochan T, Yamane I, Kashiwazaki Y. (1999). Seroprevalence of antibodies to *Neospora caninum* and associated abortion in dairy cattle from central Thailand. *Vet Parasitol.* 15;86(1):49-57.
- Wouda, W., Th.Dijkstra, A.M.H., Kramer, C. van Maanen and J.M.A. Brinkhof (1999). Seroepidemiological evidence for a relationship between *Neospora caninum* infections in dogs and cattle. *Int. J. Parasitol.* 29, 1677-1682.