

## PHYSICO - CHEMICAL CHARACTERS OF LOCALLY ISOLATED BOVINE EPHEMERAL FEVER VIRUS (BEFV)

MADBOULY, H. M \*.; HEGAZI A. G.\*\*; SOUFY H.\*\*; TAMAM S. M. \*and GAMIL G.S.\*\*

\* : Faculty of Veterinary Medicine Virology Department., Beni Suef University

\*\* : National Research Center Dokki, Gzia

Received: 27.8.2006.

Accepted: 27.8.2006

### SUMMARY

locally isolated strain of bovine ephemeral fever virus particles from outbreak in native breed 2000-2001 was examined by electron microscope and showed bullet or cone shaped, with an average length of 140-180 nm and diameter of 45-85 nm after passage (I/C) on baby mice or Vero121 and BHK21 cell line, the complete virus particles consisted of spikes, a membranous envelope and a helical nucleocapsid. This spike were easily disrupted. The harvest of BEF virus contained defective interfering particles. These particles were short cone-shaped forms about one - third the length of the infectious particles and similar in morphology to infectious virions . These particles interfering with virus replications. The BEF virus was affected by storage at 4°C and -20 °C losing its infective titer 1-1.1 log<sub>10</sub> at 4°C for 18 hours and 1.6 and 1.7 log<sub>10</sub> at -20°C for 2 months. The virus lost it's infective titer within 20 min at 56°C,

and was very sensitive to ether and chloroform as it decreased 4 and 2.6 log<sub>10</sub> when treated , respectively. The isolated BEFV was not able to agglutinate any RBCs of different species.

### INTRODUCTION

Bovine ephemeral fever (BEF) is an arthropod-borne viral disease of cattle and water buffaloes characterized by short duration, fever, stiffness, lameness, and sometimes paralysis. Both onset and recovery of infection are sudden. (St George 1988 and 1994; Nandi and Negi, 1999 Radostits et al., 2000).

The disease has a variety of names including three days sickness, stiff sickness, dengue fever of cattle, bovine epizootic fever and lazy man's disease (St. George, 1994). However, the name of bovine ephemeral fever is most commonly used. (Coml 1978; Doherty, 1978).

The causative virus was registered in the international catalogue of Arboviruses by Doherty (1978) as Bovine ephemeral fever (BEF) virus. The morbidity may be high but the mortality is low. The disease causes great economic losses such as mortalities, abortion, decreased body weight, disruption of markets, drop in milk production ranges from 34-94 with an average of 46% and cows milk yield did not reach to pre-illness level on convalescence and from lowered fertility of bulls as well as the expenses of care or treatment and vaccination (Davis et al., 1984; St-George, 1986 and 1996, Nandi and Negi, 1999, Sayed et al., 2001 and Yeruham et al., 2003) The disease was first described in Africa (Schwenk, 1867) it has been reported in many countries in Australia and Asia (St. George, 1986).

Although a viral aetiology was suspected, the cause of the disease remained unknown until van der Westhuizen, (1967) succeeded in isolating BEF virus from defibrinated blood of cattle with syndrome of ephemeral fever by intracerebral inoculation of suckling mice. He also adapted the virus to grow in cell culture (BHK - 21, clone 13). The virus was isolated from cattle and mixed with *Culicoides* spp in 1972 - 1973 (Davies and Walker, 1974).

The disease has been occurred in summer and autumn in subtropical and temperate regions of Africa, Asia and Australia. (St-George, 1984 and Nandi and Negi, 1999). It spread by insect

vectors viz., mosquitoes and *Culicoides* biologically and mechanically (Burgess, 1971, Standfast and Dyce, 1972).

The dynamics of transmission depend on the vector's abundance, distribution, host preference, and susceptibility to infection (Standfast and Dyce, 1972; Burgess, 1977; St. George 1993 and Radoszinski et al., 2000).

The disease was firstly described in Egypt by Rabagliati (1924). Since that time no publications on the occurrence of BEF in Egypt could be traced until the summer of 1991, where typical form of the disease has been recorded in different Governorates (Hassan et al., 1991; Nagi et al., 1992; and Banoub 1994). Then reoccurred of the disease at summer 2000 as severe (Ali et al., 2001; Hamoda et al., 2002, Rahman et al., 2002 and Shehabe 2004). Isolation and identification of the virus was carried out from affected animals (Hassan et al., 1991; Banoub 1994; Ali et al., 2001; Rahman et al., 2002; Shehabe 2004; Madbouly et al., 2006)

The present work aimed to study physico-chemical characters of locally isolated bovine ephemeral fever virus.

## MATERIAL AND METHODS

### 1-Virus:

Standard BEF virus (Webster's Strain) was obtained from Virology Department Faculty of Veterinary Medicine, Cairo University.

**b-Local isolated :** Isolated BEFV from native breed cattle during outbreak 2000-2001 in middle Egypt Madbouly et al., (2006)

## **2-Sera and antisera:**

### **2-1 Sera :**

**2-1-1 Foetal Calf Serum (FCS) Approved (Virus and Mycoplasma Screened).** It is purchased from GIBCO limited, Paesly, Scotland, U.K.

**2-1-2 Control negative serum.** Was kindly obtained from VSVRI Abbassia, Cairo

### **2-1-3 Reference BEF virus anti sera :**

Anti BEFV reference hyperimmune sera was obtained from Veterinary Serum and Vaccine Research Institute, (VSVRI) Abbasia, Cairo..

## **3-Experimental animals :**

### **Suckling mice :**

Albino Swiss mice 1-3 days old, obtained from the animal house of National Research Centre (NRC) these used for virus isolation and virus titration.

## **4-Tissue Culture Media :**

**4-1 Eagle's Minimum Essential Medium (MEM):** Commercially MEM (Flow laboratories, U.K)

**4-2 Growth medium Eagle's Minimum essential Medium with Earle's balanced salt solution con-**

taining 10% foetal calf serum (FCS),

## **4-3 Maintenance Medium:**

Was prepared as growth medium without foetal-calf serum (FCS) was reduce to 1-5%

## **Methods :**

**1-Preparation of hyper immune serum against BEFV:** was done According to (Davies and Walker, 1974).

**2-Propagation of the isolated virus:** the isolated virus was prepared in mice and an BHK21 or Vero121 cells according to Van-Der Westhuzen, (1967).

## **3-Titration of the isolated virus:**

### **a-Mouse adapted BEFV :**

It was done according to Theodoridis (1973).

### **b-Titration of the virus on Vero121 cells:**

It was carried out according to Theodoridis. (1973).

## **4-BEFV purification:**

It was done according to (Della - Porta and Snowdon, 1979).

## **5-Virus characterization:**

Locally isolated strain of BEF virus was studied

for effect of preservation at 4°C for 18 hours, (1985).  
 freezing at -20 °C , heat stability at 56°C according to Heushele (1970) , diethyl- ether sensitivity according to Van Der Westhuizen, (1967) and chloroform sensitivity according to Cybinski and Zakrzewski, (1983) .

### 6-Electron Microscopy :

It was performed in a guidance of Della. Part. Brown,( 1978) .The shape and size of BEFV determined by Electron microscope, with 4% monium mylbdate.

### Haemagglutination test :

It was performed according to Kaneko, et al.,

### RESULTS

**Table (1) : Effect of heat on the infectivity titer of BEFV**

Time	Heat	Log <sub>10</sub> MLD <sub>50</sub>		Log <sub>10</sub> Difference
		Before treatment	After treatment	
5 min	56°C	4.3	2.3	2
10 min	56°C	4.3	1.1	3.2
15 min	56°C	4.3	0.7	3.6
20 min	56°C	4.3	0	4.3

MLD : Mean lethal dose

Data presented in table (1) showed that the heat effect for BEF<sub>1</sub> virus at 56°C for 20 minutes. The lost its infectivity titer. While its infectivity titer decrease 2,3,2, 3,6 and 4.3 after heating at 56°C 10, 15 and 20 minutes, respectively.

**Table (2) Effect of preservation at 4°C and -20°C on the infectivity titer of BEFV.**

Type of effect	Virus	Log <sub>10</sub> MLD <sub>50</sub>		Log <sub>10</sub> difference
		Before treatment	After treatment or preservation	
preservation of BEFV <sub>1</sub> at 4°C	6 <sup>th</sup> passage of BEFV <sub>1</sub>	4.7	3.7	1
	Reference strain	5.5	4.4	1.1
preservation of BEFV <sub>1</sub> at -20°C	6 <sup>th</sup> passage of BEFV <sub>1</sub>	4.7	3	1.7
	Reference strain	5.5	3.9	1.6

MLD : Mean lethal dose

Data presented in table (2) showed clearly that the 6<sup>th</sup> passage BEF<sub>1</sub> virus was decreased one log preservation at 4°C for 18 hours. While the reference one was decreased 1.1 log<sub>10</sub> at the same time viruses were decreased 1.7 and 1.6 log<sub>10</sub> after preservation at -20°C for 2 months.

**Table (3) Effect of Diethyl ether 25% and Chloroform on the infectively titer of BEFV.**

Type of effect	Virus	Log 10 MLD 50		Log difference
		Before treatment	After treatment or preservation at 4°C	
Chloroform(5%)	BEFV local isolate	3.9	1.3	2.6
	Reference strain	4.5	2.1	2.4
Diethyl ether 25%	6 <sup>th</sup> passage of BEFV1	4.7	0.7	4
	Reference strain	5.5	1.1	4.4

Data presented in table (3) showed clearly that the BEFV and reference strain were decreased 2.6 and 2.4 log 10 after treatment with chloroform 5% for 30 minutes respectively. Both viruses were decreased 4 and 4.4 log<sub>10</sub> when treated with diethyl ether with 25% at 4°C for 18 hours.

**BEF did not agglutinated RBCs of different species.** stained, without fixation, with 4 % ammonium molybdate.

The local isolated BEFV and reference strain did not agglutinated % and 10% RBCs of human, different animal species and bird. The virus shaped was cone like shape or small bullet spike shape with helical nucleocapside coated with enveloped with short spikes as in photo (1) and (2).


**Electron microscopy BEF virus.**

This was performed as Della Porta and Brown (1979), using a EM microscope in National Research Center(NRC), Dokki, Giza. Sample were

Different shaped of virus particles were seemed in field like cone shape. Circular like shape.



(1) Electron micrographs of BEFV stained with 4% ammonium molybdate X100.000 (bullet, cone shape, Truncated cone shaped particles).



(2) Electron micrographs of BEFV stained with 4% ammonium molybdate X80.000 (bullet, cone shape, Truncated cone shaped particles).

## DISCUSSION

Bovine ephemeral fever remains as a viral disease of considerable importance to many countries including Egypt. The disease was first described by Rabagliati (1924) and this time and until summer 1991 no publication about occurrence of the disease, after that typical forms of the disease has

been occurred in different Governorate in Egypt (Hassan et al., 1991; Banoub, 1994; Sayed et al. 2001; Shehab et al., 2004).

The isolated BEF virus from native breed call during outbreak at 2000 as well as the reference one, were characterized by studying the effect of heat, Diethyl ether and chloroform, the agglutination

bility by RBCs, observation EM and purification by ultracentrifugation. The obtained result revealed that BEFV was inactivated at 56°C for 20 min table (1), sensitive to ether and chloroform table (3), not agglutinated human, duck, chicken, goose, pigeon, turkeys, buffaloes, cattle, sheep, goat, and horse RBCs. This result agrees with results agree with Kaneko et al., (1986) who reported that no hemagglutination was detected with ephemeral fever virus or another Australian virus, Berrimah (Dpp 63).

Concerning the process of purification, BEF virus on 15 to 45% sucrose density gradient ultracentrifugation, fractions of each gradient from the upper and lower of the gradient tube to obtain BEFV, this result agrees with (Della - Porta and Snowdon 1979 Dell Porta and Brown, 1980).

Concerning examination of purified BEF virus lower and upper peak in electron microscope by using 4% ammonium molybdate, it showed that lower peak contains bullet shape photo(1), cone shape photo (2) and a number of other different bullet shaped forms of virus. The upper grade contains a homogenous population of truncated cone shaped particles about one third the size of the lower grade. The cause of variation from typical bullet shape virions may be due to the temperature at which the virus grown (Della Porta and Snowdon 1979) of to the growth of interfering

virus particles Murphy et al., (1972). Or may be characteristic of BEF virus, varying slightly with different isolate like other rhabdoviruses, BEF virus produces a defective interfering (DI) particle. This DI particles is similar to the virion is being cone- shape and is about one third the size of the infectious particles similar to vesicular stomatitis virus and rabies virus DI particles (Crick et al., 1969, Crick and Brown 1974 and Della Porta and Snowdon, 1979). These DI interfere with replication of BEF virus but not interfere with VSV ( Della - Porta and Brown 1978). The BEF virus isolated strain during BEF outbreaks at 2000 and 2001 are cone or bullet shaped on morphology, that particles is similar of rabies and VSV this finding agrees with (Della Porta and Brown 1979; Della porta and Snowdon 1980).

The BEF virus shape and size bullet or cone shapes 70X145nm and slight tapered toward the rounded end. The outer envelope is closely opposed to electrons dense shell, a large particle ranged from 140-160 nm length and 85-90 nm diameter.

This finding agrees with (Van-Der Westhuizen 1967; Della Porta and Brown 1979; Della Porta and Snowdon 1980; Cheng et al., 1992 and Nandi and Negi 1999).

## REFERENCES

- Ali, N.M.; L.A. Ahmed and M.A. Shahein (2001). Isolation and identification of three day sickness virus in Egypt. *Vet. Med. J. Giza* 49,425-434
- Banoub, M. (1994). Ephemeral fever (Three days sickness) in closed milking cow from at Sharquia. 6th Sci. Cong. Faculty. *Vet. Med. Aussit* : 558-563.
- Burgess, G. W. (1971): Bovine ephemeral fever. A Review. *Vet. Bull* 41 (11) : 887-895.
- Burgess, G.W. (1977). Amicrotiter serum neutralization test for bovine ephemeral fever virus. *Australian J. Exp. Biol. Med. Sci.* 52 : 815-855.
- Cheng, L.; G. Shonlin; J. Shaode, T. D. St. George, and M. F. Uren (1992): Studies on the morphology of a Chinese isolate of bovine ephemeral fever virus. *Bovine ephemeral fever and related rhabdoviruses* proceeding of the 1st international symposium, Beijing, PRC, 25-27 ACIAR, PRC 44.
- Combs, G. P. (1978): Bovine ephemeral fever proceeding 82nd Meeting U.S. Animal Health Association, p. 29-35.
- Crick, J. and F. Brown (1974) : an interfering component of rabies virus which contains RNA. *Journal of general virology* 22, 147-151.
- Crik., J., B. Cartwright and F. Brown (1969): A study of the interference phenomenon in Vesicular stomatitis virus replication. *Archiv fur die gesamte Virusforschung* 27, 221-235.
- Cybinski, D. H. and H. Zakrzewski (1983): The isolation and preliminary characterization of a rhabdovirus in Australia related to bovine ephemeral fever virus. *Vet. Microbiol.* 8: 221-235.
- Davis, F.G.; t. Shaw and P., Ochieng. (1984). Observations on the epidemiology of ephemeral fever In Kenya. *Journal of Hygiene*, 75 : 231-235.
- Davis, G. and A.R. Walker (1974): the isolation of ephemeral fever virus from cattle and *Culicoides midges* in Kenya. *Veterinary Review*, 7 : 63-64.
- Della-Porta, A. J. and F. Brown (1978): The physico-chemical characterization of bovine ephemeral fever virus as a Member of Family Rhabdoviridae. *J. Gen. Virol.* 44:99-112.
- Della-Porta, A. J. and W.A. Snowdon (1979): An experimental inactivated virus vaccine against bovine ephemeral fever. 2. Neutralizing antibodies protect against infection. *Vet. Microbiol.*, 197-208.
- Della-Porta, A.J. and W.A. Snowdon (1980). In rhabdovirus (Bishop D.H.L., ed) chemical Rubberco Press. West-plam Beach. Florida, 3 : 167-191.
- Doherty, R. L. (1978) : "Bovine ephemeral fever. *American Journal of tropical Meicine and Hygiene* 27, 380-382.
- Hamoda, F.K.; S.S. Khalaf-Allah and M.H. Khodeir (2002). Some clinical epidemiological and laboratory studies on Bovine Ephemeral fever (three days sickness. *Vet. Med. J. Giza*, 50 : 2, 203-220.
- Hassan, H. B.; N.A. El-Danaf; M. A. M. Hafez; A. M. Ragab and M. M. Fatthia (1991): Clinico-Diagnostic studies on bovine ephemeral fever with detection of its virus for the first time in Egypt. First scientific congress; Egyptian Society for Cattle Diseases, 1-3 Dec. 1991 Assuit Egypt.
- Heuschele, W. P. (1970): Bovine ephemeral fever. I. Characteristics of the Causative virus. *Arch. Gesamte Virusforsch.*



forsch 30 : 195-202.

complication Assuit veterinary Medicine Journal 45 (92)

- Kancko, N.; Y. Inaba; H. Akashi; and K. Satou, ( 1985).  
Hemagglutination with rhabdoviruses related to bovine  
ephemeral fever virus. Veterinary - Microbiology, 11 :  
1-2-11.
- Madbouly, H.M; A.G.Hegazi, Mona El-Shabrawy,  
S.M.Tamam and .S.Gamil (2006):Isolation and identifi-  
cation of Bovine ephemerel fever virus from cattle in  
Middle Egypt (under Press)
- Murphy. F.a.; W.P. Taylor; C.A. Mims and S.G Whntfield  
(1972). Bovine ephemerel fever virus in cell culture and  
mice Archivfur die gesamte virus forschung, 38 : 234-  
249.
- Nagi, A.A.; A. I. El-Mashed; G. F. El.Bagowy and A. M.  
Moustafa (1992). An investigation on ephemerel fever  
three days sickness) in cattle at Kalyubia Governorate  
(Egypt). Annals of Agric. Sci. Moshohor, 30 : 2, 1159-  
1169.
- Nandi, S. and B.S. Negi (1999): Bovine ephemerel fever a  
review comparative immunology, Microbiology and In-  
fectious diseases. 22 : 2, 81 - 91.
- Powell, K. and D. Watason, (1975). Purification of herps  
simples virus J. Gen. Virology, 29:167-168.
- Rabagliati, D. S. (1924): Three days fever stiff sickness in  
Cattle. The Vet. Rec. 4, 23, 503-505.
- Radostits, O.M.; C.C. Gay; C.C. Blood, and K.W. Hinchel-  
iff (2000). Veterinary medicine a text book of the dis-  
ease of cattle, sheep, pigs, goats and horses. 9th Ed.  
W.B. Saunders Company Ltd. 944-960.
- Rahman, A. A., Sayed, A. A.; Sadek, A. H. and Nawa, A.  
Mohamed (2002) : "Bovine ephemerel fever Isolation of  
the causative virus and associating bacteria respiratory
- 196-212.
- Sayed, A.S.; A.H. Sadjek; and A.A. Ali (2001). Bovine  
ephemeral fever in Assiut Governorate : clinical laborato-  
ry and therapeutic studies Ausiut. Vet. Med. J. 44 (88)  
157-175.
- Schweinfurth, G. (1867): The heart of Africa. Sampson,  
Low and Searle, London, PP. 280-281.
- Shehab, G.; Abdel-Ghaffar and Rahiman I. H. A. (2004) :  
"Application of Avidin biotin complex immunoperoxi-  
dase and immunofluorescent techniques for diagnosis of  
bovine ephemerel fever Vet. Med. J.Giza. 52 : 1 : 135-  
147.
- Standfast, H.A. and A.L. Dyce (1972). Potential vectors of  
arboviruses of cattle and buffaloes in Australia Aus. Vet.  
J. 48, 224-227.
- St. George, T.D. (1984). Studies on the pathogenesis of bo-  
vine ephemerel fever in sentinel cattle. I virology & se-  
rology Australian of Biological Science 37: 341-345.
- St. George, T.D., (1986): The epidemiology of bovine  
ephemeral fever in Australia and its economic effect. In  
St. George, T.D., B.H. Kay and J. Blok.; (eds) Pproc. of  
the Fourth Symposium on Arbovirus Research in Aus-  
tralia, Brisbane Australia Canberra. Commonwealth sci-  
entific and industrial Research organization.
- St. George, T.D. (1988): Bovine ephemerel fever; A review  
trop. Animal Health Prod. 20 : 194-202.
- St. George, T.D. (1993): The natural history of ephemerel  
fever of cattle. In bovine ephemerel fever and related  
Rhabdovirus. Efs. St. George, T.D. Uren; M.F. Young;  
A.P.L. and D. Haffmer, Combera, Australian center for  
international agricultural research PP : 13.

- St. George, T.D. (1994). Bovine ephemeral fever. In infectious diseases of livestock with special reference to southern Africa, edited by JAW Coetzer, GR Thomson & RX Tustin. Capetown : oxford University press PP 404-415.
- St. George, T.D. (1995). Studies on pathogenesis of bovine ephemeral fever. *Vet. Microbial*, 46:144-148.
- St. George, T.D. (1996). Bovine ephemeral fever. In foreign animals Diseases, U.S animal Health Assoc. PP 118-128.
- Tanaka, Y., Y. Inaba; K. Sato; H.Itlo; T. Omori and M.Matsumoto (1972): Japanese journal microbial. 13, 169-176.
- Theodorides, A.; W.H. Boshoff and I.J. Dutoit (1973): Studies on the development of vaccine against bovine ephemeral fever Onderstepoort. *J. Vet. Res.*, 46 : 125-127.
- Uren, M.F., T.D. St. George and G.M. Murphy (1992). Studies on the pathogenesis of bovine ephemeral fever in experimental cattle. *Virological and Biochemical data. Vet Microbiol*, 30 : 297-307.
- Van Der Westhuizen, B. (1967): Studies on bovine ephemeral fever I. Isolation and preliminary characterization of a virus from natural and experimental produced cases of bovine ephemeral fever. *Onderstepoort J. Vet. Res.* 34 (1): 29-40.
- Yeruham, I., Braverman, Y., Yadin, H., Van Ham., Chai, D., Tiomkin, O. and Frank, D. (2003) : "Epidemiological investigation of outbreak of bovine ephemeral fever in Israel Veterinary Record 151, 117-121.